

RELATIVE MAL'TSEV CATEGORIES

TOMAS EVERAERT, JULIA GOEDECKE, TAMAR JANELIDZE-GRAY
AND TIM VAN DER LINDEN

ABSTRACT. We define relative regular Mal'tsev categories and give an overview of conditions which are equivalent to the relative Mal'tsev axiom. These include conditions on relations as well as conditions on simplicial objects. We also give various examples and counterexamples.

1. Introduction

In the paper [8], the concept of *higher-dimensional extension* and its relationship to simplicial resolutions is studied in an axiomatic setting. One of the main results [8, Theorem 3.13] relates the so-called *relative Mal'tsev axiom* to a relative Kan-property of simplicial objects. A. Carboni, G. M. Kelly and M. C. Pedicchio showed in [5] that in a regular category \mathcal{A} every simplicial object being Kan is equivalent to \mathcal{A} being a Mal'tsev category. Therefore this relative Mal'tsev axiom suggests the study of relative Mal'tsev categories. To fit into the framework of relative semi-abelian [26], relative homological [24] and relative Goursat categories [13], we now slightly alter the axioms from [8] and review some of the results of that paper in this new context. We also make a connection to properties on relative relations studied in [25], copying the ideas of [5] in this relative setting. This leads to a definition of *relative Mal'tsev categories*. Many properties of such relative Mal'tsev categories were already studied in [25], but the name did not appear there. The papers [12], [13] and [15] also mention or use this setting of relative Mal'tsev categories.

2. Axioms for extensions

The axioms we work with in this paper revolve around the concept of *higher-dimensional extension* [11, 6, 7, 8]. These higher-dimensional extensions are a particular kind of higher-dimensional arrows, which we define first.

The first author's research was supported by Fonds voor Wetenschappelijk Onderzoek (FWO-Vlaanderen). The second author's research was supported by the FNRS grant *Crédit aux chercheurs* 1.5.016.10F. The third author's research was supported by a Claude Leon Foundation postdoctoral fellowship. The fourth author works as *chargé de recherches* for Fonds de la Recherche Scientifique-FNRS. His research was supported by Fundação para a Ciência e a Tecnologia (grant number SFRH/BPD/38797/2007) and by CMUC at the University of Coimbra.

2000 Mathematics Subject Classification: 18A20, 18E10, 18G25, 18G30, 20J.

Key words and phrases: higher extension; simplicial resolution; Mal'tsev condition; relative homological algebra; arithmetical category.

© Tomas Everaert, Julia Goedecke, Tamar Janelidze-Gray and Tim Van der Linden, 2013. Permission to copy for private use granted.

The main point here is that if the pair $(\mathcal{A}, \mathcal{E})$ satisfies axioms (E1)–(E3), then so does $(\text{Ext}(\mathcal{A}), \mathcal{E}^1)$ (by [8, Proposition 1.6]; see also [11]). This allows us to iterate the definition to obtain:

2.4. DEFINITION. Given $(\mathcal{A}, \mathcal{E})$ satisfying (E1)–(E3), an n -**dimensional extension** is an n -dimensional arrow

$$\begin{array}{ccc} a_1 & \xrightarrow{f_1} & b_1 \\ a \downarrow & \Rightarrow & \downarrow b \\ a_0 & \xrightarrow{f_0} & b_0 \end{array}$$

for which the $(n-1)$ -dimensional arrows a, b, f_1 and f_0 as well as the induced arrow to the pullback of b and f_0 (in $\text{Arr}^{n-2}(\mathcal{A})$) are $(n-1)$ -dimensional extensions. We write \mathcal{E}^{n-1} for the so obtained class of n -dimensional extensions, and $\text{Ext}^n(\mathcal{A})$ for the full subcategory of $\text{Arr}^n(\mathcal{A})$ determined by the elements of \mathcal{E}^{n-1} .

The leading example for a class of extensions is the class of all regular epimorphisms in a regular category. Defining such classes of extensions axiomatically does not only give new examples, but also allows the treatment of *higher* extensions and extensions at the same time, without needing to remember which “level” is needed at any given moment—see, for instance, [8, Proposition 3.11]. Some examples of such classes of extensions can be found in [8] and also at the end of this paper.

As follows from Proposition 1.16 of [8], higher extensions can be regarded as n -cubes with certain properties (see also Remark 1.7 and Theorem 2.17 in [8]). This viewpoint ignores the specified directions which a higher arrow carries, since those are irrelevant for the extension property. Another point made in [8] is that in a precise sense, *truncated simplicial resolutions are higher-dimensional extensions*.

When the pair $(\mathcal{A}, \mathcal{E})$ satisfies additional axioms apart from (E1)–(E3) as defined above, more connections can be drawn to simplicial objects and in particular to a relative Kan property of simplicial objects. The axioms for a class of extensions \mathcal{E} in a category \mathcal{A} used in [8] for this purpose are:

- (E1) \mathcal{E} contains all isomorphisms;
- (E2) pullbacks of morphisms in \mathcal{E} exist in \mathcal{A} and are in \mathcal{E} ;
- (E3) \mathcal{E} is closed under composition;
- (E4) if $g \circ f \in \mathcal{E}$ then $g \in \mathcal{E}$ (right cancellation);
- (E5) the \mathcal{E} -**Mal'tsev axiom**: any split epimorphism of extensions

$$\begin{array}{ccc} A_1 & \begin{array}{c} \xrightarrow{f_1} \\ \xleftarrow{\quad} \end{array} & B_1 \\ a \downarrow & & \downarrow b \\ A_0 & \begin{array}{c} \xrightarrow{f_0} \\ \xleftarrow{\quad} \end{array} & B_0 \end{array}$$

in \mathcal{A} is a double extension.

Some examples in a pointed category \mathcal{A} also satisfy the stronger axiom (E5⁺) given a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ker}(a) & \longrightarrow & A_1 & \xrightarrow{a} & A_0 \longrightarrow 0 \\ & & \downarrow k & & \downarrow f & & \parallel \\ 0 & \longrightarrow & \text{Ker}(b) & \longrightarrow & B & \xrightarrow{b} & A_0 \longrightarrow 0 \end{array}$$

in \mathcal{A} with short exact rows and a and b in \mathcal{E} , if $k \in \mathcal{E}$ then also $f \in \mathcal{E}$.

2.5. REMARK. Notice that (E1) and (E4) together imply that all split epimorphisms are in \mathcal{E} and that, in a pointed category, Axiom (E2) ensures the existence of kernels of extensions. If, furthermore, \mathcal{E} consists of normal epimorphisms, then Axiom (E5⁺) implies (E5): consider a split epimorphism of extensions as in (E5). Take kernels of a and b to obtain a split epimorphism of short exact sequences:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \text{Ker}(a) & \xrightarrow{\text{ker}(a)} & A_1 & \xrightarrow{a} & A_0 & \longrightarrow & 0 \\ & & \downarrow k & \uparrow & \downarrow f_1 & \uparrow & \downarrow f_0 & \uparrow & \\ 0 & \longrightarrow & \text{Ker}(b) & \xrightarrow{\text{ker}(b)} & B_1 & \xrightarrow{b} & B_0 & \longrightarrow & 0 \end{array}$$

Since k is a split epimorphism and thus an element of \mathcal{E} , (E5⁺) implies that the right hand square is a double extension.

This axiom (E5⁺) appears in definitions of relative homological and relative semi-abelian categories (see, for instance, Condition 2(b) of Definition 1.13 of [26]).

3. Weakened axioms

As mentioned above, (E1) and (E4) imply that all split epimorphisms are in the class \mathcal{E} . However, most results of [8] can be adapted to hold in a slightly altered setting, where we assume a weak cancellation property instead of (E4) and an additional factorisation axiom.

We first give those axioms which change slightly:

(E4⁻) if $f \in \mathcal{E}$ and $g \circ f \in \mathcal{E}$ then $g \in \mathcal{E}$;

(E5⁻) any split epimorphism of extensions

$$\begin{array}{ccc} A_1 & \xrightleftharpoons{f_1} & B_1 \\ \downarrow a & & \downarrow b \\ A_0 & \xrightleftharpoons{f_0} & B_0 \end{array}$$

in \mathcal{A} with f_1 and f_0 in \mathcal{E} is a double extension.

3.1. PROPOSITION. *Let $(\mathcal{A}, \mathcal{E})$ satisfy (E1)–(E4⁻). Then \mathcal{E} contains all split epimorphisms if and only if (E4) holds.*

PROOF. By (E1), one of the implications is obvious. To prove the other, let $g \circ f$ be in \mathcal{E} . Pulling back induces the following commutative diagram:

$$\begin{array}{ccccc} P & \xrightarrow{\bar{f}} & B \times_C B & \xrightleftharpoons{\pi_1} & B \\ \uparrow \downarrow \overline{\pi_0} & \lrcorner & \uparrow \downarrow \pi_0 & \lrcorner & \downarrow g \\ A & \xrightarrow{f} & B & \xrightarrow{g} & C. \end{array}$$

The split epimorphism $\overline{\pi_0}$ is in \mathcal{E} by assumption. Furthermore, the composite $\pi_1 \circ \bar{f}$ is in \mathcal{E} by (E2). Now (E3) and (E4⁻) imply that g is in \mathcal{E} . ■

Clearly, when \mathcal{E} contains all split epimorphisms, (E5⁻) is equivalent to (E5).

These weaker axioms are satisfied, for example, by all **relative homological categories** as defined in [24]. These are pairs $(\mathcal{A}, \mathcal{E})$, where \mathcal{A} is a pointed category with finite limits and cokernels, and \mathcal{E} is a class of normal epimorphisms in \mathcal{A} satisfying axioms (E1)–(E3), (E4⁻) and (E5⁺), as well as the axiom

- (F) if a morphism f in \mathcal{A} factors as $f = e \circ m$ with m a monomorphism and $e \in \mathcal{E}$, then it also factors (essentially uniquely) as $f = m' \circ e'$ with m' a monomorphism and $e' \in \mathcal{E}$.

Using (F), several results of [8] about higher extensions go through, and we obtain new examples. In particular, we have the following lemma.

3.2. LEMMA. *If \mathcal{A} has finite products, \mathcal{E} is a class of epimorphisms in \mathcal{A} and $(\mathcal{A}, \mathcal{E})$ satisfies (E1)–(E3) and (F), then given any split epimorphism of extensions*

$$\begin{array}{ccccc} A_1 \times_{A_0} A_1 & \rightrightarrows & A_1 & \xrightarrow{a} & A_0 \\ r \downarrow \uparrow & & f_1 \downarrow \uparrow & & f_0 \downarrow \uparrow \\ B_1 \times_{B_0} B_1 & \rightrightarrows & B_1 & \xrightarrow{b} & B_0 \end{array}$$

with f_1 and f_0 in \mathcal{E} , taking kernel pairs of a and b gives an extension r .

PROOF. Consider a diagram as above and the composite morphism

$$A_1 \times_{A_0} A_1 \xrightarrow{\langle \pi_0, \pi_1 \rangle} A_1 \times A_1 \xrightarrow{f_1 \times f_1} B_1 \times B_1.$$

The product $f_1 \times f_1$ is an extension by (E2) and (E3), and $\langle \pi_0, \pi_1 \rangle$ is a monomorphism. Hence by (F) the morphism $(f_1 \times f_1) \circ \langle \pi_0, \pi_1 \rangle$ admits a factorisation $\langle r_0, r_1 \rangle \circ e$, where (R, r_0, r_1) is a relation on B_1 and e is in \mathcal{E} . Since e is an epimorphism by assumption, we have $b \circ r_0 = b \circ r_1$, and R is contained in $B_1 \times_{B_0} B_1$. Now r being a split epimorphism implies that $R = B_1 \times_{B_0} B_1$. ■

In [8, Proposition 3.3] Axiom (E5) is shown to be equivalent to three other conditions involving double extensions, namely Axiom (E4) holding for the class of double extensions \mathcal{E}^1 , every split epimorphism of split epimorphisms being a double extension and a condition relating the comparison map between the kernel pairs of two morphisms in a square to the property of the square being a double extension. In the present setting, the axiom (E5⁻) is not quite equivalent to all the corresponding conditions. Instead we have:

3.3. PROPOSITION. *Let $(\mathcal{A}, \mathcal{E})$ satisfy (E1)–(E3) and (E4⁻). Consider the following statements:*

- (i) (E4⁻) holds for \mathcal{E}^1 , that is, if $g \circ f \in \mathcal{E}^1$ and $f \in \mathcal{E}^1$ then $g \in \mathcal{E}^1$;
- (ii) Axiom (E5⁻) holds;
- (iii) every split epimorphism of split epimorphisms with a, b, f_1 and f_0 in \mathcal{E} , i.e. every diagram

$$\begin{array}{ccc} A_1 & \xleftarrow{\bar{f}_1} & B_1 \\ \uparrow a & \bar{a} & \downarrow b \\ & f_1 & \uparrow \bar{b} \\ A_0 & \xleftarrow{\bar{f}_0} & B_0 \\ & f_0 & \end{array}$$

such that $f_0 a = b f_1$, $\bar{f}_0 \bar{b} = a \bar{f}_1$, $\bar{b} f_0 = f_1 \bar{a}$, $\bar{a} \bar{f}_0 = \bar{f}_1 \bar{b}$ and $f_0 \bar{f}_0 = 1_{B_0}$, $f_1 \bar{f}_1 = 1_{B_1}$, $a \bar{a} = 1_{A_0}$, $b \bar{b} = 1_{B_0}$ and the four split epimorphisms are in \mathcal{E} , is a double extension;

- (iv) given a diagram

$$\begin{array}{ccccc} A_1 \times_{B_1} A_1 & \rightrightarrows & A_1 & \xrightarrow{f_1} & B_1 \\ \downarrow r & & \downarrow a & & \downarrow b \\ A_0 \times_{B_0} A_0 & \rightrightarrows & A_0 & \xrightarrow{f_0} & B_0 \end{array}$$

in \mathcal{A} with a, b, f_1 and f_0 in \mathcal{E} , the arrow r is in \mathcal{E} if and only if the right hand side square is in \mathcal{E}^1 .

Then (ii) \Rightarrow (iv) \Rightarrow (i) and (ii) \Rightarrow (iii). If $(\mathcal{A}, \mathcal{E})$ also satisfies (F), then (iii) \Rightarrow (iv) \Rightarrow (ii), resulting in the equivalence of (ii), (iii) and (iv).

PROOF. Because of Lemma 3.2, the proofs of the implications (ii) \Rightarrow (iii) \Rightarrow (iv) \Rightarrow (i) are very similar to those found in [8, Proposition 3.3]. For (ii) \Rightarrow (iv) a modified version of the proof of [8, Lemma 3.2] may be used, and (iv) \Rightarrow (ii) is straightforward. ■

It can be seen that (E1)–(E4⁻) and (E5⁻) “go up to higher dimensions together”, meaning:

3.4. PROPOSITION. *Let \mathcal{A} be a category and \mathcal{E} a class of arrows in \mathcal{A} . If $(\mathcal{A}, \mathcal{E})$ satisfies (E1)–(E4⁻) and (E5⁻), then $(\text{Ext}(\mathcal{A}), \mathcal{E}^1)$ satisfies the same conditions.*

PROOF. The proof of [8, Proposition 3.4] can easily be adapted to this weaker situation. ■

4. The relative Mal'tsev axiom and relations

Classically, Mal'tsev categories are defined using properties of relations. Therefore we now connect the relative Mal'tsev condition $(E5^-)$ to the conditions on \mathcal{E} -relations studied in [26, 25]. For this, we use a context given in Condition 2.1 in [26], that is, we assume that \mathcal{A} has finite products, \mathcal{E} is a class of regular epimorphisms in \mathcal{A} and $(\mathcal{A}, \mathcal{E})$ satisfies axioms $(E1)$ – $(E3)$, $(E4^-)$ and (F) . In [13] such a pair $(\mathcal{A}, \mathcal{E})$ is called a **relative regular category**. For a more detailed explanation see [26] and [13].

4.1. DEFINITION. Given two objects A and B in \mathcal{A} , an \mathcal{E} -relation from A to B is a subobject of $A \times B$ such that for any representing monomorphism $\langle r_0, r_1 \rangle: R \rightarrow A \times B$, the morphisms $r_0: R \rightarrow A$ and $r_1: R \rightarrow B$ are in \mathcal{E} .

Using the axioms given, such \mathcal{E} -relations can be composed and this composition is associative. The usual definitions and calculations of relations apply. This setting allows us to copy proofs and methods from [5] to a relative situation. Many of these results were proved in [25, Theorem 2.3.6]; in particular, for a relative regular category $(\mathcal{A}, \mathcal{E})$, we have:

4.2. PROPOSITION. For any relative regular category $(\mathcal{A}, \mathcal{E})$, the following are equivalent:

- (i) for equivalence \mathcal{E} -relations R and S on an object A in \mathcal{A} , the relation $SR: A \rightarrow A$ is an equivalence \mathcal{E} -relation;
- (ii) any two equivalence \mathcal{E} -relations R and S on an object A in \mathcal{A} permute: $SR = RS$;
- (iii) any two \mathcal{E} -effective equivalence relations R and S (i.e., kernel pairs of extensions) on A in \mathcal{A} permute;
- (iv) every \mathcal{E} -relation is difunctional;
- (v) every reflexive \mathcal{E} -relation is an equivalence \mathcal{E} -relation;
- (vi) every reflexive \mathcal{E} -relation is symmetric;
- (vii) every reflexive \mathcal{E} -relation is transitive. ■

This suggests a definition of a relative regular Mal'tsev category to fit into the context of relative homological and relative semi-abelian categories.

4.3. DEFINITION. A relative regular category $(\mathcal{A}, \mathcal{E})$ is **relative Mal'tsev** if it satisfies any one of the conditions 4.2(i)–4.2(vii) above.

In their paper [15], M. Gran and D. Rodelo showed that the axiom $(E5^-)$ is also equivalent to several other conditions, including a condition on relations and a diagram lemma called the *Relative Cuboid Lemma*:

4.4. THEOREM. [15] *If $(\mathcal{A}, \mathcal{E})$ is a relative regular category, then the following are equivalent:*

- (i) *Axiom $(E5^-)$;*
- (ii) *any two \mathcal{E} -effective equivalence relations R and S on A in \mathcal{A} permute;*
- (iii) *for any commutative cube*

$$\begin{array}{ccccc}
 W \times_D C & \xrightarrow{v} & Y \times_B A & & \\
 \uparrow \downarrow & \searrow & \uparrow \downarrow & \searrow & \\
 & & C & \xrightarrow{c} & A \\
 \downarrow & \swarrow & \downarrow & \swarrow & \downarrow \\
 W & \xrightarrow{g} & Y & \xrightarrow{f} & B \\
 \downarrow & \searrow & \downarrow & \searrow & \\
 & & D & \xrightarrow{d} & B
 \end{array}$$

in \mathcal{A} , where f and g are split epimorphisms in \mathcal{E} , c , d , and w are in \mathcal{E} , and the left and right squares are pullbacks, the induced morphism $v: W \times_D C \rightarrow Y \times_B A$ is an extension;

(iv) *the Relative Split Cuboid Lemma holds;*

(v) *the Relative Upper Cuboid Lemma holds.* ■

Notice that any relative regular Mal'tsev category is **relative Goursat** in the sense of [13]: for equivalence \mathcal{E} -relations R and S on an object A , the equality $RSR = SRS$ holds. Hence in any relative regular Mal'tsev category, also the *Relative 3×3 Lemma* is valid—see [28, 16, 13].

We are now ready to extend a main result about the relative Mal'tsev axiom from [8]—its characterisation in terms of the \mathcal{E} -Kan property for \mathcal{E} -simplicial objects—to the current setting (Proposition 4.7).

4.5. DEFINITION. Let \mathbb{A} be a simplicial object and consider $n \geq 2$ and $0 \leq k \leq n$. The **object of (n, k) -horns in \mathbb{A}** is an object $A(n, k)$ together with arrows $a_i: A(n, k) \rightarrow A_{n-1}$ for $i \in \{0, \dots, n\} \setminus \{k\}$ satisfying

$$\partial_i \circ a_j = \partial_{j-1} \circ a_i \text{ for all } i < j \text{ with } i, j \neq k$$

which is universal with respect to this property. We also define $A(1, 0) = A(1, 1) = A_0$.

A simplicial object is **\mathcal{E} -Kan** when all $A(n, k)$ exist and all comparison morphisms $A_n \rightarrow A(n, k)$ are in \mathcal{E} . In particular, the comparison morphisms to the $(1, k)$ -horns are $\partial_0: A_1 \rightarrow A(1, 0)$ and $\partial_1: A_1 \rightarrow A(1, 1)$.

In fact, we can still use the proof of Proposition 3.12 in [8], but we need to show that certain split epimorphisms are in \mathcal{E} . We shall again need:

4.6. PROPOSITION. *In a relative regular category $(\mathcal{A}, \mathcal{E})$, an augmented \mathcal{E} -simplicial object \mathbb{A} which is contractible and \mathcal{E} -Kan is always an \mathcal{E} -resolution: for all $n \geq -1$, the factorisation $A_{n+1} \rightarrow K_{n+1}\mathbb{A}$ to the simplicial kernel $K_{n+1}\mathbb{A}$ of $\partial_0, \dots, \partial_n: A_n \rightarrow A_{n-1}$ (and $K_0\mathbb{A} = A_{-1}$) is in \mathcal{E} .*

PROOF. The proof of Proposition 3.9 in [8] goes through, if we can show that the morphism r in the diagram

$$\begin{array}{ccccccc}
 A_{n+2} & \longrightarrow & A(n+2, 0) & \xrightarrow{a_1} & A_{n+1} & \xrightarrow{\partial_1} & A_n \\
 \downarrow \partial_0 & & \uparrow r & \xrightarrow{a_{n+2}} & \downarrow \partial_0 & \xrightarrow{\partial_{n+1}} & \downarrow \partial_0 \\
 A_{n+1} & \xrightarrow{\langle \partial_0, \dots, \partial_{n+1} \rangle} & K_{n+1}\mathbb{A} & \xrightarrow{k_0} & A_n & \xrightarrow{\partial_0} & A_{n-1} \\
 & & & \xrightarrow{k_{n+1}} & & \xrightarrow{\partial_n} &
 \end{array}$$

is an extension. This is done as in the proof of Lemma 3.2. Note that the simplicial kernels $K_{n+1}\mathbb{A}$ exist for this \mathcal{E} -resolution, since [8, Lemma 3.8] uses only axioms (E1)–(E3). ■

4.7. PROPOSITION. *Let $(\mathcal{A}, \mathcal{E})$ be a relative regular category such that \mathcal{A} has simplicial kernels. Then (E5⁻) holds if and only if every \mathcal{E} -simplicial object in \mathcal{A} is \mathcal{E} -Kan.*

PROOF. The proofs of Propositions 3.11 and 3.12 from [8] may easily be adapted to this setting, however for the implication \Leftarrow some additional steps are needed.

When (E1)–(E4⁻) and (F) hold and every \mathcal{E} -simplicial object is \mathcal{E} -Kan, we wish to show that every split epimorphism of split epimorphisms with all appropriate arrows in \mathcal{E} is a double extension. This then implies (E5⁻) by Proposition 3.3. We can first reduce the situation to a (truncated) contractible augmented \mathcal{E} -simplicial object

$$\begin{array}{ccc}
 & \xleftarrow{\sigma_{-1}} & \\
 & \downarrow \partial_0 & \\
 A_1 & \xleftarrow{\sigma_0} & A_0 \xrightarrow{\partial_0} A_{-1} \\
 & \downarrow \partial_1 & \\
 & \xleftarrow{\sigma_1} &
 \end{array}
 \tag{A}$$

Given a split epimorphism of split epimorphisms

$$\begin{array}{ccc}
 A & \xleftarrow{\bar{f}} & B \\
 \uparrow a & \xrightarrow{f} & \uparrow b \\
 A' & \xleftarrow{\bar{f}'} & B' \\
 \downarrow a & \xrightarrow{f'} & \downarrow b
 \end{array}$$

with a, b, \bar{f} and \bar{f}' in \mathcal{E} , we define $A_{-1} = B'$, $A_0 = A$, $\partial_0 = f' \circ a = b \circ f: A_0 \rightarrow A_{-1}$ and $\sigma_{-1} = \bar{a} \circ \bar{f}' = \bar{f} \circ \bar{b}: A_{-1} \rightarrow A_0$. The morphisms ∂_0 and $\partial_1: A_1 \rightarrow A_0$ are defined by the

pullback

$$\begin{array}{ccc}
 A_1 & \xrightarrow{p} & A_0 \\
 \downarrow \langle \partial_0, \partial_1 \rangle & \lrcorner & \downarrow \langle a, f \rangle \\
 A_0 \times_{A_{-1}} A_0 & \xrightarrow{a \times_{1_{B'}} f} & A' \times_{B'} B
 \end{array} \tag{B}$$

where the morphism $a \times_{1_{B'}} f$ is an extension as the pullback of the double extensions $(f' \circ a, f'): a \rightarrow 1_{B'}$ and $(f' \circ a, b): f \rightarrow 1_{B'}$. The morphisms $\sigma_{-1}, \sigma_0: A_0 \rightarrow A_1$ are universally induced by

$$(a \times_{1_{B'}} f) \circ \langle 1_{A_0}, 1_{A_0} \rangle = \langle a, f \rangle \circ 1_{A_0}$$

and

$$(a \times_{1_{B'}} f) \circ \langle 1_{A_0}, \bar{a} \circ \bar{f}' \circ f' \circ a \rangle = \langle a, f \rangle \circ (\bar{a} \circ a)$$

respectively. In contrast to the proof in [8], we also need $\sigma_1: A_0 \rightarrow A_1$ induced by

$$(a \times_{1_{B'}} f) \circ \langle \bar{f} \circ \bar{b} \circ b \circ f, 1_{A_0} \rangle = \langle a, f \rangle \circ (\bar{f} \circ f).$$

These morphisms then satisfy the simplicial identities; in particular, $\partial_1 \circ \sigma_1 = 1_{A_0}$ and $\partial_0 \circ \sigma_1 = \sigma_{-1} \circ \partial_0$. It remains to check that ∂_0 and ∂_1 are also extensions. We may decompose the diagram defining, say, ∂_0 , as

$$\begin{array}{ccccc}
 A_1 & \xrightarrow{\bar{r}} & Q & \longrightarrow & A_0 \\
 \downarrow \langle \partial_0, \partial_1 \rangle & \lrcorner & \downarrow \langle a, f \rangle & \lrcorner & \downarrow \langle a, f \rangle \\
 A_0 \times_{A_{-1}} A_0 & \xrightarrow{r} & P & \longrightarrow & A' \times_{B'} B \\
 \downarrow \pi_0 & & \downarrow \bar{\pi}_{A'} & \lrcorner & \downarrow \pi_{A'} \\
 A_0 & \xlongequal{\quad} & A_0 & \xrightarrow{a} & A'
 \end{array}$$

The induced morphism r is an extension (since the bottom rectangle is a double extension), hence so is \bar{r} . The composite $\bar{\pi}_{A'} \circ \langle a, f \rangle$ is also an extension, as a pullback of $a = \pi_{A'} \circ \langle a, f \rangle$. Hence $\partial_0 = \pi_0 \circ \langle \partial_0, \partial_1 \rangle$ is an extension by (E3). Similarly, so is ∂_1 .

A truncated \mathcal{E} -simplicial object of the shape (\mathbf{A}) can be extended to a contractible augmented simplicial object \mathbb{A} by constructing successive simplicial kernels. Using (F) we now show that such a simplicial object is actually an \mathcal{E} -simplicial object, so that it is \mathcal{E} -Kan by assumption. To see this, we write (\mathbf{A}) in the form of a cube, where A_2 is the induced simplicial kernel. The simplicial identities ensure that all possible squares in it

- Axiom (E5⁻) holds;
- every \mathcal{E} -simplicial object is \mathcal{E} -Kan. ■

Theorem 4.8 tells us that a relative Mal'tsev category is exactly a pair $(\mathcal{A}, \mathcal{E})$ satisfying axioms (E1)–(E3), (E4⁻), (E5⁻) and (F).

5. On the axiom (F)

We explain under which conditions, in the absolute case, Axiom (F) goes up to higher dimensions. Here \mathcal{E} is the class of all regular epimorphisms in \mathcal{A} .

5.1. REMARK. Notice that a morphism $f = (f_1, f_0): a \rightarrow b$ between extensions a and b is a monomorphism in $\text{Ext}(\mathcal{A})$ if and only if f_1 is a monomorphism. In particular, there are no restrictions on f_0 . When \mathcal{A} is regular, pushouts of regular epimorphisms are exactly the regular epimorphisms in $\text{Ext}(\mathcal{A})$.

5.2. PROPOSITION. *Let \mathcal{A} be a regular category and \mathcal{E} the class of all regular epimorphisms in \mathcal{A} . The following conditions are equivalent:*

- (i) \mathcal{A} is exact Mal'tsev;
- (ii) the pushout of an extension by an extension exists and is a double extension;
- (iii) $(\text{Ext}(\mathcal{A}), \mathcal{E}^1)$ satisfies (F).

PROOF. The equivalence of (i) and (ii) was proved by A. Carboni, G. M. Kelly and M. C. Pedicchio in [5]. Assuming (ii), any morphism $f: a \rightarrow b$ in $\text{Ext}(\mathcal{A})$ factors as a double extension followed by a monomorphism as follows.

$$\begin{array}{ccccc}
 A_1 & \xrightarrow{e} & I & \xrightarrow{m} & B_1 \\
 \downarrow a & \Rightarrow & \downarrow & \Rightarrow & \downarrow b \\
 A_0 & \longrightarrow & P & \longrightarrow & B_0
 \end{array}$$

Here $f_1 = m \circ e$ is the regular epi-mono factorisation of f_1 and the left hand square is the pushout of e by a . Note that the former exists because \mathcal{A} is regular and the latter by assumption. Hence, (ii) implies (iii). To see that (iii) implies (ii), consider extensions f and g and the morphism of extensions

$$\begin{array}{ccc}
 A & \xrightarrow{f} & B \\
 \downarrow g & \Rightarrow & \downarrow \\
 C & \longrightarrow & 1
 \end{array}$$

where 1 is the terminal object. This square can be factored as a monomorphism (in the category of extensions) followed by a double extension as follows.

$$\begin{array}{ccccc} A & \xlongequal{\quad} & A & \xrightarrow{f} & B \\ g \downarrow & \Rightarrow & \downarrow & \Rightarrow & \downarrow \\ C & \longrightarrow & 1 & \xlongequal{\quad} & 1 \end{array}$$

The assumption implies that the square can also be factored as a double extension followed by a monomorphism.

$$\begin{array}{ccccc} A & \xrightarrow{e} & I & \xrightarrow{m} & B \\ g \downarrow & \Rightarrow & \downarrow & \Rightarrow & \downarrow b \\ C & \longrightarrow & I' & \longrightarrow & 1 \end{array}$$

But this means in particular that m is a monomorphism. Hence, it is an isomorphism, since it is also a regular epimorphism (as f is). It follows that the pushout of f by g exists (it is given by the left hand square) and is a double extension, as desired. ■

Let us now investigate under which circumstances (F) “goes up” to $(\text{Ext}^2(\mathcal{A}), \mathcal{E}^2)$. Clearly, as soon as $(\text{Ext}^2(\mathcal{A}), \mathcal{E}^2)$ satisfies (F), the same will be true for $(\text{Ext}(\mathcal{A}), \mathcal{E}^1)$. Hence, by Proposition 5.2, a necessary condition for $(\text{Ext}^2(\mathcal{A}), \mathcal{E}^2)$ to satisfy (F) is that \mathcal{A} is exact Mal’tsev. Observe that, in this case, $\text{Ext}(\mathcal{A})$ is regular: regular epimorphisms in $\text{Ext}(\mathcal{A})$ are double extensions, which we know are pullback-stable. Hence, we can apply Proposition 5.2 to $\text{Ext}(\mathcal{A})$ and find, in particular, that the pair $(\text{Ext}^2(\mathcal{A}), \mathcal{E}^2)$ satisfies (F) if and only if $\text{Ext}(\mathcal{A})$ is exact Mal’tsev.

Now, recall from [29] that an exact Mal’tsev category is **arithmetical** if every internal groupoid is an equivalence relation. Examples of arithmetical categories are the dual of the category of pointed sets, more generally, the dual of the category of pointed objects in any topos, and also the categories of von Neumann regular rings, Boolean rings and Heyting semi-lattices. It was proved in [2] that an exact Mal’tsev category is arithmetical if and only if the category $\text{Equiv}(\mathcal{A})$ of internal equivalence relations in \mathcal{A} is exact. In this case $\text{Equiv}(\mathcal{A})$ is in fact again arithmetical and, in particular, exact Mal’tsev. Since, moreover, there is a category equivalence $\text{Equiv}(\mathcal{A}) \simeq \text{Ext}(\mathcal{A})$ because \mathcal{A} is exact, we have:

5.3. PROPOSITION. *Let \mathcal{A} be an exact Mal’tsev category and \mathcal{E} the class of all regular epimorphisms in \mathcal{A} . The following are equivalent:*

- \mathcal{A} is arithmetical;
- $\text{Ext}(\mathcal{A})$ is arithmetical;
- $\text{Ext}(\mathcal{A})$ is exact Mal’tsev;
- any pushout of a double extension by a double extension exists (in the category $\text{Ext}(\mathcal{A})$) and is a three-fold extension;
- $(\text{Ext}^2(\mathcal{A}), \mathcal{E}^2)$ satisfies (F). ■

5.4. **REMARK.** Notice that Proposition 5.3 also implies that Axiom (F) is satisfied by $(\text{Ext}^n(\mathcal{A}), \mathcal{E}^n)$ for *every* n as soon as the category \mathcal{A} is arithmetical. Conversely, the category \mathcal{A} is arithmetical as soon as there exists an $n \geq 2$ such that (F) holds for $(\text{Ext}^n(\mathcal{A}), \mathcal{E}^n)$.

Since being arithmetical is a rather restrictive property for a (Mal'tsev) category to have, we can conclude this analysis by saying that Axiom (F) "hardly ever" goes up to $(\text{Ext}^2(\mathcal{A}), \mathcal{E}^2)$ or higher.

6. Examples

We end this article with several examples and counterexamples. Some of the examples satisfy the stronger axiom $(E5^+)$, cf. [1, 6, 7, 24].

6.1. **EXAMPLE.** [Relative homological categories] Under the axioms $(E1)$ – $(E4^-)$ and (F), Axiom $(E5^+)$ also implies $(E5^-)$ as soon as \mathcal{E} is a class of epimorphisms. Hence relative homological and relative semi-abelian categories as defined in [24, 26] are relatively Mal'tsev, but generally need not satisfy the stronger $(E4)$ and $(E5)$. An example of a relative semi-abelian category is a semi-abelian category \mathcal{A} with \mathcal{E} being the class of central extensions in the sense of Huq, closed under composition [25, Proposition 5.3.2]; see also Example 6.4. That is, any morphism in \mathcal{E} is the composition of regular epimorphisms $f: A \rightarrow B$ with $[\text{Ker}(f), A] = 0$, where $[\text{Ker}(f), A]$ is the commutator of $\text{Ker}(f)$ and A in the sense of Huq [18].

When \mathcal{E} is a class of regular epimorphisms in a regular Mal'tsev category \mathcal{A} satisfying $(E1)$ – $(E2)$, then it is easy to check that $(E3)$, $(E4^-)$ and $(E5^-)$ hold as soon as the following **two out of three property** is satisfied: given a composite $g \circ f$ of regular epimorphisms $f: A \rightarrow B$ and $g: B \rightarrow C$, if any two of $g \circ f$, f and g lie in \mathcal{E} , then so does the third. We shall make use of this fact when considering the following two examples, which are given by categorical Galois theory [19, 20]. Notice that this uses the regular Mal'tsev property to show that, in the square given in $(E5^-)$, the comparison to the pullback is already a regular epimorphism, and then the two out of three property shows that it is in fact in \mathcal{E} .

6.2. **EXAMPLE.** [Trivial extensions] Let \mathcal{B} be a full and replete reflective subcategory of a regular Mal'tsev category \mathcal{A} . Write $H: \mathcal{B} \rightarrow \mathcal{A}$ for the inclusion functor and $I: \mathcal{A} \rightarrow \mathcal{B}$ for its left adjoint. Assume that HI preserves regular epimorphisms and I is *admissible* [20] with respect to regular epimorphisms. This means that I preserves all pullbacks of the form

$$\begin{array}{ccc}
 B \times_{HI(B)} H(X) & \longrightarrow & H(X) \\
 \downarrow & \lrcorner & \downarrow_{H(\varphi)} \\
 B & \xrightarrow{\eta_B} & HI(B)
 \end{array} \tag{C}$$

where $\varphi: X \rightarrow I(B)$ is a regular epimorphism. For instance, \mathcal{B} could be a Birkhoff subcategory of \mathcal{A} (a full reflective subcategory closed under subobjects and regular quotients) if \mathcal{A} is also Barr-exact (see [21]).

Recall that a **trivial covering** or **trivial extension** (with respect to I) is a regular epimorphism f such that the commutative square induced by the unit $\eta: 1_{\mathcal{A}} \Rightarrow HI$

$$\begin{array}{ccc} A & \xrightarrow{\eta_A} & HI(A) \\ f \downarrow & & \downarrow HI(f) \\ B & \xrightarrow{\eta_B} & HI(B) \end{array} \quad (\mathbf{D})$$

is a pullback. With \mathcal{E} the class of all trivial extensions, $(\mathcal{A}, \mathcal{E})$ satisfies conditions (E1)–(E4⁻) and (E5⁻); see also [27]. (The stronger axiom (E4) need not hold as in general not every split epimorphism is a trivial extension: for instance, when \mathcal{A} is pointed, a morphism $A \rightarrow 0$ is a trivial extension if and only if A is in \mathcal{B} .) Indeed, the validity of (E1) is clear while (E2) follows from the admissibility of I (see Proposition 2.4 in [22]). Hence, it suffices to prove the two out of three property, of which only one implication is not immediate. To see that $g: B \rightarrow C$ is a trivial extension as soon as $f: A \rightarrow B$ and $g \circ f$ are, it suffices to note that, since $HI(f)$ is a pullback-stable regular epimorphism, the change of base functor $(HI(f))^*: (\mathcal{A} \downarrow HI(B)) \rightarrow (\mathcal{A} \downarrow HI(A))$ is conservative [23].

When \mathcal{A} is Barr-exact and \mathcal{B} is a Birkhoff subcategory of \mathcal{A} , then $(\mathcal{A}, \mathcal{E})$ also satisfies (F). Indeed, condition (F) is easily inferred from the fact that in this case the square (D) is a pushout, hence a regular pushout (a double extension) for any regular epimorphism f [5, 21]. If moreover \mathcal{A} is pointed with cokernels and \mathcal{B} is protomodular, then $(\mathcal{A}, \mathcal{E})$ forms a relative homological category [27].

6.3. EXAMPLE. [Torsion theories] Recall that $p: E \rightarrow B$ is an **effective descent morphism** if the change of base functor $p^*: (\mathcal{A} \downarrow B) \rightarrow (\mathcal{A} \downarrow E)$ is monadic. Let \mathcal{A} be a homological category in which every regular epimorphism is effective for descent (for instance, \mathcal{A} could be semi-abelian) and let \mathcal{B} be a torsion-free subcategory of \mathcal{A} (a full regular epi-reflective subcategory of \mathcal{A} such that the associated radical $T: \mathcal{A} \rightarrow \mathcal{A}$ is idempotent, see [4]). Then the reflector $I: \mathcal{A} \rightarrow \mathcal{B}$ is semi-left exact: it preserves all pullbacks of the form (C), now for *all* morphisms $\varphi: X \rightarrow I(B)$. In particular, the previous example applies. Thus we find that the pair $(\mathcal{A}, \mathcal{E})$ satisfies conditions (E1)–(E4⁻) and (E5⁻), for \mathcal{E} the class of all trivial extensions.

Let us now write \mathcal{E}^* for the class of (regular epi)morphisms $f: A \rightarrow B$ that are “locally in \mathcal{E} ”, in the sense that there exists an effective descent morphism $p: E \rightarrow B$ in \mathcal{A} such that the pullback $p^*(f): E \times_B A \rightarrow E$ is in \mathcal{E} . The morphisms in \mathcal{E}^* are usually called **coverings** or **central extensions**. While the pair $(\mathcal{A}, \mathcal{E}^*)$ satisfies conditions (E1) and (E2) because $(\mathcal{A}, \mathcal{E})$ does, \mathcal{E}^* is in general not closed under composition. However, it was shown in [10] that \mathcal{E}^* is composition-closed as soon as the reflector I is **protoad-ditive** [9, 10]: I preserves split short exact sequences. Let us briefly recall the argument. First of all, it was shown in [10] that the central extensions with respect to I (which we

shall, from now on, assume to be protoadditive) are exactly those regular epimorphisms $f: A \rightarrow B$ whose kernel $\text{Ker}(f)$ is in \mathcal{B} . Now, let $f: A \rightarrow B$ and $g: B \rightarrow C$ be regular epimorphisms. Then we have a short exact sequence

$$0 \longrightarrow \text{Ker}(f) \longrightarrow \text{Ker}(g \circ f) \longrightarrow \text{Ker}(g) \longrightarrow 0$$

and we see that $g \circ f$ is a central extension as soon as f and g are, since the torsion-free subcategory \mathcal{B} is closed under extensions (which means that when $\text{Ker}(f) \in \mathcal{B}$ and $\text{Ker}(g) \in \mathcal{B}$ then $\text{Ker}(g \circ f) \in \mathcal{B}$) [4]. Furthermore, since \mathcal{B} is a (regular epi)-reflective subcategory of \mathcal{A} , \mathcal{B} is closed under subobjects, and so f is a central extension as soon as $g \circ f$ is. If we assume that \mathcal{B} is, moreover, closed under regular quotients (which is equivalent to \mathcal{B} being a Birkhoff subcategory of \mathcal{A}) then g is a central extension as soon as $g \circ f$ is, and we may conclude that \mathcal{E}^* satisfies the two out of three property. Once again using that \mathcal{B} is closed under subjects in \mathcal{A} , it is easily verified that the pair $(\mathcal{A}, \mathcal{E}^*)$ also satisfies Axiom (F). (Note that the same two out of three property can be used to show that $(\mathcal{A}, \mathcal{E}^*)$ is, in fact, relatively homological.)

Examples of such an \mathcal{A} and \mathcal{B} are given, for instance, by taking \mathcal{A} to be the category of compact Hausdorff groups and \mathcal{B} the subcategory of profinite groups [10], or \mathcal{A} to be the category of internal groupoids in a semi-abelian category and \mathcal{B} the subcategory of discrete groupoids [9]. Since a reflector into an epi-reflective subcategory of an abelian category is necessarily (proto)additive, any cohereditary torsion theory (meaning that \mathcal{B} is closed under quotients) in an abelian category \mathcal{A} provides an example as well. However, there are no non-trivial examples in the categories of groups or of abelian groups, as follows from Proposition 5.5 in [30].

6.4. EXAMPLE. [Composites of central extensions] We use the context of Example 6.2, assuming in addition that \mathcal{A} is Barr-exact and \mathcal{B} is a Birkhoff subcategory of \mathcal{A} . In this setting a regular epimorphism $f: A \rightarrow B$ is a central extension (with respect to I) if there exists a regular epimorphism $p: E \rightarrow B$ such that the pullback $p^*(f): E \times_B A \rightarrow E$ of f along p is a trivial extension. We take \mathcal{E} to be the class of composites of such central extensions. If now \mathcal{A} is pointed and has cokernels and coproducts, and \mathcal{B} is protomodular, then $(\mathcal{A}, \mathcal{E})$ forms a relative semi-abelian category [27]. When \mathcal{B} is determined by the abelian objects in \mathcal{A} , we regain the example mentioned in 6.1: then the \mathcal{B} -central extensions in \mathcal{A} are determined by the Smith commutator [3], while, via [17], extensions are Smith-central if and only if they are Huq-central as in Example 6.1.

6.5. EXAMPLE. [Internal groupoids] Let the pair $(\mathcal{A}, \mathcal{E})$ satisfy axioms (E1)–(E4⁻), (E5⁻) and (F). Denote by $\mathbf{Gpd}_{\mathcal{E}}(\mathcal{A})$ the category of **internal \mathcal{E} -groupoids** in \mathcal{A} : groupoids G in \mathcal{A} with the property that all split epimorphisms occurring in the diagram of G are in \mathcal{E} . Write $\overline{\mathcal{E}}$ for the class of degree-wise \mathcal{E} -extensions. Then $(\mathbf{Gpd}_{\mathcal{E}}(\mathcal{A}), \overline{\mathcal{E}})$ is relatively Mal'tsev. Indeed, to see that axioms (E2) and (E5⁻) are satisfied, observe that pullbacks along morphisms in $\overline{\mathcal{E}}$ are degree-wise pullbacks in \mathcal{A} . For Axiom (F) note that products are computed degree-wise as well, and that $\mathbf{Gpd}_{\mathcal{E}}(\mathcal{A})$ is closed in $\mathbf{RG}_{\mathcal{E}}(\mathcal{A})$ —the category of “reflexive \mathcal{E} -graphs” in \mathcal{A} —under “ $\overline{\mathcal{E}}$ -quotients”, as a consequence of the relative Mal'tsev condition for $(\mathcal{A}, \mathcal{E})$. See [14] for the absolute case.

6.6. EXAMPLE. [Regular pullback squares] This is an example of a pair $(\mathcal{A}, \mathcal{E})$ which satisfies (E1)–(E4⁻) and (E5⁻), but where not every split epimorphism is an extension, nor does (F) hold. We take \mathcal{A} to be the category $\mathbf{ExtGp}_{\text{tf}}$ of extensions (regular epimorphisms) in the category of torsion-free groups. The class \mathcal{E} consists of regular pullback squares, i.e., pullbacks of regular epimorphisms. It is easy to find a split epimorphism of extensions which is not a pullback, and it is also easy to see that (E1)–(E4⁻) and (E5⁻) hold using that \mathbf{Gp}_{tf} is regular Mal'tsev. We give a counterexample for Axiom (F); it is based on the fact that pushouts in \mathbf{Gp}_{tf} are different from pushouts in \mathbf{Gp} and may not be regular pushouts. They are constructed by reflecting the pushout in \mathbf{Gp} into the subcategory \mathbf{Gp}_{tf} .

An example of a pushout in \mathbf{Gp}_{tf} which is not a pushout in \mathbf{Gp} is the square

$$\begin{array}{ccc} \mathbb{Z} \times_{\mathbb{Z}_2} \mathbb{Z} & \longrightarrow & \mathbb{Z} \\ \downarrow & & \downarrow \\ \mathbb{Z} & \longrightarrow & 0. \end{array} \quad (\mathbf{G})$$

(\mathbb{Z}_2 is torsion while \mathbb{Z} is torsion-free.) The diagram

$$\begin{array}{ccccc} \mathbb{Z} \times_{\mathbb{Z}_2} \mathbb{Z} & \longrightarrow & \mathbb{Z} \times \mathbb{Z} & \longrightarrow & \mathbb{Z} \\ \downarrow & & \downarrow \lrcorner & & \downarrow \\ \mathbb{Z} & \xlongequal{\quad} & \mathbb{Z} & \longrightarrow & 0 \end{array}$$

now displays a monomorphism composed with an \mathcal{E} -extension which cannot be written as an \mathcal{E} -extension composed with a monomorphism, as the square (\mathbf{G}) is not in \mathcal{E} .

References

- [1] D. Bourn, *3 × 3 Lemma and protomodularity*, J. Algebra **236** (2001), 778–795.
- [2] D. Bourn, *A categorical genealogy for the congruence distributive property*, Theory Appl. Categ. **8** (2001), no. 14, 391–407.
- [3] D. Bourn and M. Gran, *Central extensions in semi-abelian categories*, J. Pure Appl. Algebra **175** (2002), 31–44.
- [4] D. Bourn and M. Gran, *Torsion theories in homological categories*, J. Algebra **305** (2006), 18–47.
- [5] A. Carboni, G. M. Kelly, and M. C. Pedicchio, *Some remarks on Maltsev and Goursat categories*, Appl. Categ. Structures **1** (1993), 385–421.
- [6] T. Everaert, *An approach to non-abelian homology based on Categorical Galois Theory*, Ph.D. thesis, Vrije Universiteit Brussel, 2007.
- [7] T. Everaert, *Higher central extensions and Hopf formulae*, J. Algebra **324** (2010), 1771–1789.

- [8] T. Everaert, J. Goedecke, and T. Van der Linden, *Resolutions, higher extensions and the relative Mal'tsev axiom*, J. Algebra **371** (2012), 132–155.
- [9] T. Everaert and M. Gran, *Homology of n -fold groupoids*, Theory Appl. Categ. **23** (2010), no. 2, 22–41.
- [10] T. Everaert and M. Gran, *Protoadditive functors, derived torsion theories and homology*, preprint arXiv:1111.5448v1, 2011.
- [11] T. Everaert, M. Gran, and T. Van der Linden, *Higher Hopf formulae for homology via Galois Theory*, Adv. Math. **217** (2008), no. 5, 2231–2267.
- [12] J. Goedecke, *Homology in relative semi-abelian categories*, Appl. Categ. Structures, in press, 2012.
- [13] J. Goedecke and T. Janelidze, *Relative Goursat categories*, J. Pure Appl. Algebra **216** (2012), 1726–1733.
- [14] M. Gran, *Internal categories in Mal'cev categories*, J. Pure Appl. Algebra **143** (1999), 221–229.
- [15] M. Gran and D. Rodelo, *The cuboid lemma and Mal'tsev categories*, Pré-Publicações DMUC **12-32** (2012), 1–15, submitted.
- [16] M. Gran and D. Rodelo, *A new characterisation of Goursat categories*, Appl. Categ. Structures **20** (2012), no. 3, 229–238.
- [17] M. Gran and T. Van der Linden, *On the second cohomology group in semi-abelian categories*, J. Pure Appl. Algebra **212** (2008), 636–651.
- [18] S. A. Huq, *Commutator, nilpotency and solvability in categories*, Quart. J. Math. Oxford **19** (1968), no. 2, 363–389.
- [19] G. Janelidze, *Pure Galois theory in categories*, J. Algebra **132** (1990), no. 2, 270–286.
- [20] G. Janelidze, *Categorical Galois theory: revision and some recent developments*, Galois connections and applications, Math. Appl., vol. 565, Kluwer Acad. Publ., 2004, pp. 139–171.
- [21] G. Janelidze and G. M. Kelly, *Galois theory and a general notion of central extension*, J. Pure Appl. Algebra **97** (1994), no. 2, 135–161.
- [22] G. Janelidze and G. M. Kelly, *The reflectiveness of covering morphisms in algebra and geometry*, Theory Appl. Categ. **3** (1997), no. 6, 132–159.

- [23] G. Janelidze, M. Sobral, and W. Tholen, *Beyond Barr exactness: Effective descent morphisms*, *Categorical Foundations: Special Topics in Order, Topology, Algebra and Sheaf Theory* (M. C. Pedicchio and W. Tholen, eds.), *Encyclopedia of Math. Appl.*, vol. 97, Cambridge Univ. Press, 2004, pp. 359–405.
- [24] T. Janelidze, *Relative homological categories*, *J. Homotopy Relat. Struct.* **1** (2006), no. 1, 185–194.
- [25] T. Janelidze, *Foundation of relative non-abelian homological algebra*, Ph.D. thesis, University of Cape Town, 2009.
- [26] T. Janelidze, *Relative semi-abelian categories*, *Appl. Categ. Structures* **17** (2009), 373–386.
- [27] T. Janelidze-Gray, *Composites of central extensions form a relative semi-abelian category*, submitted, 2012.
- [28] S. Lack, *The 3-by-3 lemma for regular Goursat categories*, *Homology, Homotopy Appl.* **6** (2004), no. 1, 1–3.
- [29] M. C. Pedicchio, *Arithmetical categories and commutator theory*, *Appl. Categ. Structures* **4** (1996), no. 2–3, 297–305.
- [30] J. Rosický and W. Tholen, *Factorisation, fibration and torsion*, *J. Homotopy Relat. Struct.* **2** (2007), no. 2, 295–314.

Vakgroep Wiskunde, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium

Institut de recherche en mathématique et physique, Université catholique de Louvain, chemin du cyclotron 2 bte L7.01.02, 1348 Louvain-la-Neuve, Belgium

Queens' College, University of Cambridge, United Kingdom

Department of Mathematical Sciences, University of South Africa, Pretoria PO Box 392, UNISA 0003, South Africa

CMUC, University of Coimbra, 3001-454 Coimbra, Portugal

Email: `teveraer@vub.ac.be`
`julia.goedecke@cantab.net`
`tjanel@gmail.com`
`tim.vanderlinden@uclouvain.be`